

Quiver relations and associated symmetric polynomials

Dimitry Noshchenko

January 2, 2017

Abstract

The idea is to identify certain path algebra elements with symmetric functions. We propose such a morphism by solving the quiver relations, which describe the Plucker-type embedding for quiver grassmannians.

1 Introduction

Quiver is a directed graph with multiply edges and loops allowed. Let Q = finite quiver with k vertices and l arrows. Quiver representation M is a collection of vector spaces and linear maps, attached to each vertex and arrow of Q :

$$M = \{V_1, \dots, V_k\} \cup \{M_{v,i,j}\}, \quad \dim(V_i) = d_i, \quad M_{v,i,j} = \text{Mat}(d_i \times d_j, \mathbf{k}) : V_i \rightarrow V_j \quad (1)$$

Quiver grassmannian $Gr_{\underline{e}}(Q, M)$ of dimension vector $\underline{e} = (e_1, \dots, e_k)$ contains all $\underline{e}$ -dimensional subspaces of M , compatible with the underlying linear maps.

Fix an ordered basis $\mathcal{B} = \{1, 2, \dots, d_1, d_1 + 1, \dots, d_l\}$, $I \subset \mathcal{B}$ for all V_i s. Quiver relations (QR) are of the form [4]:

$$\sum_{s', t'} (-1)^{\epsilon(s', t')} M_{v, s', t'} \Delta_s^{s'} \Delta_{t'}^t - \sum_{s'} (-1)^{\epsilon(s', t)} M_{v, s', t} \Delta_s^{s'} \Delta_I = 0, \quad I \subset \mathcal{B}, \quad |I| = e \quad (2)$$

Define $Fset$ as a collection of all non-vanishing Δ_j^i 's in 2. Our aim is to generate polynomials which are sums of GL characters with coefficients given by the embedding

$$Gr_{\bar{\tau}(e)}(Q, M) \xrightarrow{\phi_1} Gr(e, d) \xrightarrow{\phi_2} \mathbb{P}^{e+d-1}, \quad e = \sum e_i, d = \sum d_i \quad (3)$$

Define 2 kind of invariants:

$$P(Q, s, \mathbf{M}) = \sum_R \Delta_R \chi_R(\bar{t}) \quad (4)$$

with Δ = homogeneous coordinate on the image of embedding 3, and

$$\tilde{P}(Q, s, \mathbf{M}) = \sum_{R'} \tilde{\Delta}_{R'} \chi_R(\bar{t}) \quad (5)$$

with $\tilde{\Delta} \in Fset$ only (PR are not taken into part). Note that P, P' are inhomogeneous in R, R' (may contain Young diagrams of different size).

Proposition 1. *In case of $\chi_\lambda = s_\lambda$ polynomials $P(Q, s, \mathbf{M})$ form a basis of Symm algebra. This seems*

to be obvious, but not the only one possible choice of χ .
Write $\mathbf{M}$ as block matrix:

$$\mathbf{M} = \begin{bmatrix} M_{1,1} & M_{1,2} & M_{1,3} & \dots \\ M_{2,1} & M_{2,2} & M_{2,3} & \\ M_{3,1} & M_{3,2} & M_{3,3} & \\ \vdots & & & \ddots \end{bmatrix} \quad (6)$$

with each $M_{i,j}$ corresponding to (i, j) linear map.

2 Computation of $P, \tilde{P}$. Quiver relations as vanishing minors

2.1 $Q = A_2, M = \{\mathbb{C}, \mathbb{C}, \mu = \text{Mat}(2, 2, \mathbb{C})\}, \bar{e} = (1, 1)$

$I = \{2, 4\}$: One non-trivial equation, $\#(+, -) = (2, 2)$

$$F(v, 2, 3) = \mu_{v,1,4} \Delta_{\{1,4\}} \Delta_{\{2,3\}} + \mu_{v,2,4} \Delta_{\{2,4\}} \Delta_{\{2,3\}} - \mu_{v,1,3} \Delta_{\{1,4\}} \Delta_{\{2,4\}} - \mu_{v,2,3} \Delta_{\{2,4\}}^2 = 0 \quad (7)$$

Or:

$$F(v, 2, 3) = \begin{vmatrix} \mu_{v,1,4} & 0 & \Delta_{\{2,4\}} \\ \mu_{v,2,4} & \Delta_{\{1,4\}} & \frac{\mu_{v,2,3} \Delta_{\{2,4\}}^2}{\mu_{v,1,4} \Delta_{\{2,3\}}} \\ \mu_{v,1,3} & \Delta_{\{2,3\}} & \Delta_{\{2,3\}} \end{vmatrix} = \begin{vmatrix} \mu_{v,1,4} & \Delta_{\{2,4\}} & \Delta_{\{2,4\}} \\ -\mu_{v,2,4} & \Delta_{\{1,4\}} & -\frac{\mu_{v,2,3} \Delta_{\{2,4\}}}{\mu_{v,1,3}} \\ \mu_{v,1,3} & 0 & \Delta_{\{2,3\}} \end{vmatrix} = 0 \quad (8)$$

$$, \quad \Delta_{1,2} = 0, \quad \Delta_{3,4} = 0 \quad (9)$$

Note that if we inverse the arrow, our equation will change:

$$F(v, 4, 1) = \mu'_{v,3,2} \Delta_{\{2,3\}} \Delta_{\{1,4\}} + \mu'_{v,4,2} \Delta_{\{2,4\}} \Delta_{\{1,4\}} - \mu'_{v,3,1} \Delta_{\{2,3\}} \Delta_{\{2,4\}} - \mu'_{v,4,1} \Delta_{\{2,4\}}^2 = 0, \quad (10)$$

where μ' is the inverse map. Now fix $\mu = \begin{pmatrix} p & 1-p \\ 1-p & p \end{pmatrix}$ - quantum permutation matrix from <https://arxiv.org/abs/1109.4888>. Combining this with the only Plu"cker identity

$$\Delta_{\{1,2\}} \Delta_{\{3,4\}} - \Delta_{\{1,3\}} \Delta_{\{2,4\}} + \Delta_{\{1,4\}} \Delta_{\{2,3\}} = 0 \quad (11)$$

we obtain our polynomial invariant:

$$\boxed{P(A_2, 4, \mu) = Qh_1 - \frac{Q(-Wp - Q + pQ)h_2}{W - Wp + pQ} + W(h_1^2 - h_2h_0) - \frac{(-Wp - Q + pQ)W(h_1h_2 - h_3h_0)}{W - Wp + pQ}} \quad (12)$$

where Q, W are arbitrary complex numbers.

2.2 $M = \{\mathbb{C}^2, \mathbb{C}^2, \mu = \text{Mat}(4, 4, \mathbb{C})\}, \bar{e} = (2, 2)$

$I = 3, 4, 7, 8$: We have 2x2 non-trivial quiver equations () with $\#(+, -) = (4, 5), (6, 3)$ One of them ()

$$F(v, 3, 5) = \begin{vmatrix} \mu_{v,1,7} + \mu_{v,2,7} + \mu_{v,2,8} & \Delta_{\{3,4,7,8\}} & \Delta_{\{3,4,5,7\}} + 2\Delta_{\{3,4,7,8\}} \\ \mu_{v,3,8} - \mu_{v,3,7} - \mu_{v,1,5} & \Delta_{\{1,4,7,8\}} + 2\Delta_{\{2,4,7,8\}} & 0 \\ -\mu_{v,1,8} + \mu_{v,2,5} + \frac{\mu_{v,3,5} \Delta_{\{3,4,7,8\}}}{\Delta_{\{2,4,7,8\}}} & \Delta_{\{3,4,7,8\}} + \Delta_{\{1,4,7,8\}} & 2\Delta_{\{3,4,5,8\}} + \Delta_{\{3,4,5,7\}} \end{vmatrix} = 0 \quad (13)$$

$$\begin{aligned} & \mu_{v,2,8}\Delta_{\{2,4,7,8\}}\Delta_{\{3,4,5,7\}} + \mu_{v,3,8}\Delta_{\{3,4,7,8\}}\Delta_{\{3,4,5,7\}} - \mu_{v,1,5}\Delta_{\{1,4,7,8\}}\Delta_{\{3,4,7,8\}} \\ & - \mu_{v,2,5}\Delta_{\{2,4,7,8\}}\Delta_{\{3,4,7,8\}} - \mu_{v,3,5}\Delta_{\{3,4,7,8\}}^2 \end{aligned}$$

Can we rewrite them in a determinantal form?

cont.

$$\begin{aligned} F(v, 4, 5) = & \begin{vmatrix} \mu_{v,4,7} & m_{2;1,2} & \Delta_{\{3,4,5,7\}} \\ 0 & \Delta_{\{3,4,7,8\}} & 0 \\ -\mu_{v,4,8} & m_{2;3,2} & \Delta_{\{3,4,5,8\}} \end{vmatrix} + \begin{vmatrix} \mu_{v,2,5} & 0 & \Delta_{\{3,4,5,8\}} \\ -\mu_{v,1,7} & \Delta_{\{2,3,7,8\}} & 0 \\ \mu_{v,2,7} & \Delta_{\{1,3,7,8\}} & \Delta_{\{3,4,7,8\}} \end{vmatrix} + \\ & + \begin{vmatrix} -\mu_{v,1,8} & \Delta_{\{2,3,7,8\}} & \Delta_{\{3,4,7,8\}} \\ \mu_{v,2,8} & \Delta_{\{1,3,7,8\}} & 0 \\ -\mu_{v,1,5} & -\frac{\mu_{v,4,5}\Delta_{\{3,4,7,8\}}}{\mu_{v,2,8}} & \Delta_{\{3,4,5,7\}} \end{vmatrix} = 0. \end{aligned} \quad (14)$$

Polynomial $P(A_2, 4, Id)$ define $P(Q, n, M)$ = linear combination of GL characters with coefficients running through the Fset. **this turns out to be a special base of Symm.** This case gives

$$Fset = \begin{pmatrix} \Delta_{\{1,4,7,8\}} & \Delta_{\{1,3,7,8\}} & & & & & & \\ \Delta_{\{2,4,7,8\}} & \Delta_{\{2,3,7,8\}} & & & & & & \\ \Delta_{\{3,4,7,8\}} & & & & & & & \\ & \Delta_{\{3,4,7,8\}} & & & & & & \\ & & \Delta_{\{1,4,7,8\}} & -\Delta_{\{1,3,7,8\}} & & & & \\ & & \Delta_{\{2,4,7,8\}} & -\Delta_{\{2,3,7,8\}} & & & & \\ & & \Delta_{\{3,4,7,8\}} & & & & & \\ & & & \Delta_{\{3,4,7,8\}} & & & & \end{pmatrix} \quad (15)$$

Now we can write down the explicit form of $P(A_2, 4, Id)$:

$$Fset = [\Delta_{\{1,3,7,8\}}, \Delta_{\{1,4,7,8\}}, \Delta_{\{2,3,7,8\}}, \Delta_{\{2,4,7,8\}}, \Delta_{\{3,4,5,7\}}, \Delta_{\{3,4,5,8\}}, \Delta_{\{3,4,6,7\}}, \Delta_{\{3,4,6,8\}}, \Delta_{\{3,4,7,8\}}] \quad (16)$$

Assuming all free parameters = ± 1 , we get the following solution:

$$Fset = [1, 1, 1, -1, -1, 1, -1, -1, 1] \quad (17)$$

Now choose $\mu = \mu(p, q)$. Solving QR's independently, we get:

$$\Delta_{\{1,3,7,8\}} = -\frac{1}{pq}(\Delta_{\{3,4,5,7\}} - 2\Delta_{\{3,4,5,7\}}q - \Delta_{\{1,4,7,8\}}p - \Delta_{\{2,4,7,8\}} + p\Delta_{\{2,4,7,8\}} + \quad (18)$$

$$+ \Delta_{\{1,4,7,8\}}pq + \Delta_{\{2,4,7,8\}}q - p\Delta_{\{2,4,7,8\}}q - \Delta_{\{2,3,7,8\}}q + p\Delta_{\{2,3,7,8\}}q)$$

$$\Delta_{\{3,4,5,8\}} = -\frac{1}{q}(\Delta_{\{3,4,5,7\}} - \Delta_{\{3,4,5,7\}}q - \Delta_{\{1,4,7,8\}}p - \Delta_{\{2,4,7,8\}} + p\Delta_{\{2,4,7,8\}}) \quad (19)$$

$$\Delta_{\{3,4,6,7\}} = -\frac{1}{p(2q-1)}(2p\Delta_{\{2,3,7,8\}}q + 2\Delta_{\{3,4,5,7\}}pq - 2p\Delta_{\{2,4,7,8\}}q - p\Delta_{\{3,4,5,7\}} + 2p\Delta_{\{2,4,7,8\}} - \quad (20)$$

$$- 2\Delta_{\{3,4,5,7\}}q + \Delta_{\{2,4,7,8\}}q - \Delta_{\{2,3,7,8\}}q + \Delta_{\{3,4,5,7\}} - \Delta_{\{2,4,7,8\}})$$

$$\Delta_{\{3,4,6,8\}} = \frac{1}{pq(2q-1)}(\Delta_{\{3,4,6,8\}} = 2p\Delta_{\{2,4,7,8\}} - 2\Delta_{\{3,4,5,7\}}q^2p - 2\Delta_{\{1,4,7,8\}}p^2q + 2p^2\Delta_{\{2,4,7,8\}}q + 2\Delta_{\{2,4,7,8\}}q^2p - \quad (21)$$

Assuming $\{\Delta_{\{1,4,7,8\}}, \Delta_{\{2,3,7,8\}}, \Delta_{\{2,4,7,8\}}, \Delta_{\{3,4,5,7\}}, \Delta_{\{3,4,7,8\}}\} = \{1\}$, we can try to resolve all remaining (Plu"cker) relations. In this case

Now we're almost ready to write our target polynomial. The only thing left is to convert the indices $i_1, i_2, \dots, i_s$ to the corresponding Young diagram. In order to do this, expand the following determinant by Cauchy-Binet formula: We have:

2.3 $Q = A_2, d_i = 6, e_i = 3, \mu = \sigma(p, q, 6)$

$$I = \left\{ \underbrace{4, 5, 6}, \underbrace{10, 11, 12} \right\} \quad (24)$$

Recall that each Δ is a $(6, 6)$ complex minor. The following minors are supposed to vanish:

$$\{\Delta_{\{1,4,5,6,10,11\}}, \Delta_{\{1,4,5,6,10,12\}}, \Delta_{\{1,4,5,6,11,12\}}, \Delta_{\{2,4,5,6,10,11\}}, \Delta_{\{2,4,5,6,10,12\}}, \Delta_{\{2,4,5,6,11,12\}}, \Delta_{\{3,4,5,6,10,11\}}, \Delta_{\{3,4,5,6,10,12\}}, \Delta_{\{3,4,5,6,11,12\}}\} = \{0\} \quad (25)$$

Total $\#QR = 3^3$. (3 for each $in\{4, 5, 6\}$: $F(v, i, j)$). For arbitrary μ these relations look really huge!

The choice $\mu = Id(6)$ kills most of the $Fset$ entries, so we can write $Fset$ as $(12, 12)$ sparse matrix with $\{0, \pm 1\}$ entries of the shape (empty space is filled with zeroes):

```
Fset=Matrix([seq(seq(Fset[1+i .. 12+i], i = 12*j), j = 0 .. 11)])
```

4

Finally, convert all survived $Fset$ entries to corresponding Young diagrams to get out target polynomial:

$$\tilde{P}(A_2, 6, Id) = s \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array} - s \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array} + s \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array} + s \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array} - s \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array} + s \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \square \\ \hline \end{array} + \dots, \quad (27)$$

total $\#(0, +, -) = (120, 21, 3)$

2.4 $Q = A_3, d_i = 2, \mu_{i=1..2} = Mat(2, 2, \mathbb{C}), \bar{e} = (1, 1, 1)\}$

Now let's add one more vertex:

$$Q = A_3 : \quad \circ \xleftarrow{\mu_1} \circ \xleftarrow{\mu_2} \circ, \quad \mu_i \quad (28)$$

$$Gr_{(1,1,1)}(A_3, M) = \quad (29)$$

Choose $I = \{2, 4, 6\}$, so now we have the embedding

$$Gr_{(1,1,1)}(A_3, M) \longrightarrow Gr(3, 6) \longrightarrow \mathbb{P}^{6+3-1} \quad (30)$$

For $v = 1$ we have the following non-trivial quiver relations:

$$F(1, 2, 3) = \mu_{1,1,4} \Delta_{\{1,4,6\}} \Delta_{\{2,3,6\}} + \mu_{1,2,4} \Delta_{\{2,4,6\}} \Delta_{\{2,3,6\}} - \mu_{1,1,3} \Delta_{\{1,4,6\}} \Delta_{\{2,4,6\}} - \mu_{1,2,3} \Delta_{\{2,4,6\}}^2 = 0 \quad (31)$$

Note that if $\mu = Id$, the solution looks as follows:

$$Fset = \begin{pmatrix} \Delta_{\{1,4,6\}} & \Delta_{\{1,2,6\}} & \Delta_{\{1,2,4\}} \\ 1 & & \\ & \Delta_{\{1,4,6\}} & \\ & 1 & \\ & & \Delta_{\{1,4,6\}} \\ & & 1 \end{pmatrix}, \quad (32)$$

where all empty cells are zeroes.

2.5 $Q = A_3, d_i = 4, \mu_1 = \sigma(p, q, 4), \mu_2 = Id$

Initial data:

$$\mathcal{B} = \{1, 2, \dots, 12\}, I = \{3, 4, 7, 8, 11, 12\} \quad (33)$$

$$\mu_1 = \begin{pmatrix} p & 1-p & 0 & 0 \\ 1-p & p & 0 & 0 \\ 0 & 0 & q & 1-q \\ 0 & 0 & 1-q & q \end{pmatrix}, \mu_2 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \quad (34)$$

This case reads

$$Fset' = \left\{ 1, \Delta_{\{2,3,4,7,8,11\}}, \Delta_{\{2,3,4,7,8,12\}}, \Delta_{\{3,4,7,8,9,11\}}, \Delta_{\{3,4,7,8,10,11\}}, -\frac{-q+2qp}{q(-1+2p)}, -\frac{2p^2q-qp}{qp(-1+2p)} \right\} \quad (35)$$

3 Diagrammatic notation and exact formulas for invariants

$\tilde{P}(A_n, 2k, \sigma)$:

$$\begin{aligned} \tilde{P}(A_2, 4, \sigma) = & \frac{(-2q - qp)(h_3^3 h_2 - h_3^2 h_1 h_4 - 2h_3 h_4 h_2^2 + 2h_4^2 h_2 h_1 - h_4 h_5 h_1^2 + h_5 h_2^3 - h_6 h_2^2 h_1 + h_6 h_3 h_1^2)}{qp} + \\ & + h_3^2 h_2^2 - h_5 h_3 h_1^2 - h_4 h_2^3 + 2h_5 h_2^2 h_1 - h_6 h_2 h_1^2 - 2h_3 h_4 h_2 h_1 + 2h_4^2 h_1^2 + h_3^2 h_2 h_1 - h_3 h_4 h_1^2 + \\ & \frac{(-2qp - 2q + p)(h_5 h_3 h_1^2 - h_6 h_2 h_1^2)}{p(2q - 1)} + \frac{(-2q + 3qp - 2q^2 p + 2q^2)(h_3 h_4 h_1^2 - h_5 h_2 h_1^2)}{pq(2q - 1)} - \\ & - h_4 h_2^2 h_1 + h_5 h_2 h_1^2 + h_4 h_5 h_1^2 - h_6 h_3 h_1^2 + h_3^2 h_1^2 - h_4 h_2 h_1^2 \end{aligned} \quad (36)$$

What is the asymptotic formula for $\#+, -$ in $Fset(Id, m)$, when $m \rightarrow \infty$? The first few terms are shown on the figure: $Fset$ matrix arrangement for A_4 quiver with $d_i = 4, \mu_i = Id$:




Figure 1: Convergence of $\{\frac{\#+}{\#-}\}$ sequence, A_2 quiver, $\mu = Id, \dim(M) = i$

$$Fset = \begin{pmatrix} x_1 & x_2 & & x_3 & x_4 & & x_5 & x_6 & & x_7 & x_8 \\ x_1 & x_{11} & & x_{12} & x_{13} & & x_{14} & x_{15} & & x_{16} & x_{17} \\ 1 & & & & & & & & & & \\ & 1 & & & & & & & & & \\ & & & x_1 & -x_2 & & x_{18} & x_{19} & & x_2 & x_{21} \\ & & & x_1 & -x_{11} & & x_{22} & x_{23} & & x_{24} & x_{25} \\ & & & 1 & & & & & & & \\ & & & & 1 & & & & & & \\ & & & & & & x_1 & x_2 & & & \\ & & & & & & x_1 & x_{11} & & & \\ & & & & & & 1 & & & & \\ & & & & & & & 1 & & & \\ & & & & & & & & & x_1 & -x_2 \\ & & & & & & & & & x_1 & -x_{11} \\ & & & & & & & & & 1 & \\ & & & & & & & & & & 1 \end{pmatrix} \quad (37)$$


where $x_j = \Delta_{I(j)}$. Compare this to $A_2, d_i = 8, \mu_i = Id$ case:

$$\left(\begin{array}{cccccccccccccccc} & x_1 & -x_2 & x_3 & -x_4 & & & & & & & & & & \\ & x_5 & x_6 & x_7 & x_8 & & & & & & & & & & \\ & x_9 & x_{10} & x_{11} & x_{12} & & & & & & & & & & \\ x_{13} & x_{14} & x_{15} & x_{16} & & & & & & & & & & & \\ & 1 & & & & & & & & & & & & & \\ & & 1 & & & & & & & & & & & & \\ & & & 1 & & & & & & & & & & & \\ & & & & 1 & & & & & & & & & & \\ & & & & & x_1 & x_2 & x_3 & x_4 & & & & & & \\ & & & & & x_5 & -x_6 & x_7 & -x_8 & & & & & & \\ & & & & & x_9 & -x_{10} & x_{11} & -x_{12} & & & & & & \\ & & & & & x_{13} & -x_{14} & x_{15} & -x_{16} & & & & & & \\ & & & & & 1 & & & & & & & & & \\ & & & & & & 1 & & & & & & & & \\ & & & & & & & 1 & & & & & & & \\ & & & & & & & & 1 & & & & & & \\ & & & & & & & & & 1 & & & & & \end{array} \right) \quad (38)$$

Totally positive *Fset* arrangement for $A_4 \times$ mutations, $d_i = 6, \mu = Id$:

[illegible]

[illegible]

 $\frac{1}{2}(\tilde{P}_+ - \tilde{P}_-)$ for cyclic 6-vertex graph:

$$\left(\begin{array}{c} -s \\ \text{Diagram 1} \end{array} \right) + \left(\begin{array}{c} +s \\ \text{Diagram 2} \end{array} \right) + \left(\begin{array}{c} \Delta \\ \text{Diagram 3} \end{array} \right) + \left(\begin{array}{c} +s \\ \text{Diagram 4} \end{array} \right) + \left(\begin{array}{c} -s \\ \text{Diagram 5} \end{array} \right) + \left(\begin{array}{c} \Delta \\ \text{Diagram 6} \end{array} \right) + \left(\begin{array}{c} +s \\ \text{Diagram 7} \end{array} \right) + \left(\begin{array}{c} -s \\ \text{Diagram 8} \end{array} \right) + \left(\begin{array}{c} +s \\ \text{Diagram 9} \end{array} \right) + \left(\begin{array}{c} \Delta \\ \text{Diagram 10} \end{array} \right)$$


Lemma 1. *Asymptotic formula for $\tilde{P}_{\max}(Q_{\text{univ}}(n), n, Id)$:*

- $m \times m$ “cubic ladder” decomposition $\simeq \Delta$ -summands in $\tilde{P}_I$
- each Δ -summand consists of Young diagrams factorized by the “ladder-block” addition

The formula:

$$\tilde{P}_{\max}(Q(m), n, Id) = \sum_{i=1..m \cdot \text{cl}(\frac{m}{2})} \sum_{j=1..n} s_{\lambda(i,j)} \Delta_{\lambda(i,*)}, \quad (42)$$

$\frac{1}{2}(\tilde{P}_+ - \tilde{P}_-)$ for cyclic 8-vertex graph:



5 Re-evaluation of $\tilde{P}$ for Coxeter quivers. “Vertex at ∞ ”

Let Q be a quiver with m vertices and all possible arrows (but only 2 opposite arrows for each pair of vertices is allowed, so the direction $v = \pm 1$). Choose $n, e \in \mathbb{N}$ and the integer-valued vectors

$$B = \{1, \dots, m \cdot n\}, I = \{jn - i + 1\}_{i=1..[n/2], j=1..m} \subset B, |I| = e, \quad (44)$$

where $[x]$ is the integer part of x . Fix the representation $R(Q)$: vertices $\sim V_i(\mathbb{Z}^n)$, arrows $\sim$ transposition matrices $M_{v,i,j} = \sigma(\lambda_{i,j})^{\text{sign}(v)}$.

Each $\sigma(\lambda_{i,j})$ is a $(n \times n)$ transposition matrix, partitioned by $\lambda_{i,j}$ for the (i,j) -th arrow. We fix *the particular choice* $R = \Sigma_0$:

$$(i, j) \rightarrow \lambda_{i,j} : (1, 4) \rightarrow \begin{array}{|c|c|c|c|} \hline \square & & & \\ \hline \square & \square & & \\ \hline \square & \square & \square & \\ \hline \square & \square & \square & \square \\ \hline \end{array}, (1, 3) \rightarrow \begin{array}{|c|c|c|} \hline \square & & \\ \hline \square & \square & \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array}, (1, 2) \rightarrow \begin{array}{|c|c|} \hline \square & \\ \hline \square & \square \\ \hline \square & \square \\ \hline \square & \square \\ \hline \end{array}, (2, 4) \rightarrow \begin{array}{|c|c|c|c|} \hline \square & & & \\ \hline \square & \square & & \\ \hline \square & \square & \square & \\ \hline \square & \square & \square & \square \\ \hline \end{array}, (2, 3) \rightarrow \begin{array}{|c|c|c|} \hline \square & & \\ \hline \square & \square & \\ \hline \square & \square & \square \\ \hline \square & \square & \square \\ \hline \end{array}, \dots \quad (45)$$

Assume also $\dim(\text{Gr}(Q, R)) = \bar{e} = \left\lfloor \frac{1}{2}n \right\rfloor$ for the corresponding quiver grassmannian [2, 4], $e_i = e_j \forall i, j$.

For instance, the case $(m, n) = (4, 4)$ is given by

$$\mathbf{M}_+ = \begin{pmatrix} 0 & \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \end{bmatrix} & \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix} & \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \\ 0 & 0 & \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{bmatrix} & \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \\ 0 & 0 & 0 & \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \\ 0 & 0 & 0 & 0 \end{pmatrix} \quad (46)$$

and $\mathbf{M}_-$ equals by-block inverse of $\mathbf{M}_+$. Choose some path $x \in \mathcal{P}(Q)$ and solve the quiver relations with vanishing conditions [4]:

$$\sum_{s', t'} (-1)_{\epsilon(s', t')} M_{v, s', t'} \Delta_s^{s'} \Delta_{t'}^t - \sum_{s'} (-1)_{\epsilon(s', t)} M_{v, s', t} \Delta_s^{s'} \Delta_I = 0, \quad (47)$$

$\Delta_J = 0$ for some non-proper J 's

Define symmetric polynomials

$$\tilde{P}_R(x) = \sum_{i, j} \Delta_{\mu_{i, j}} s_{\mu_{i, j}}(p_1, p_2, \dots), \quad (48)$$

where $\mu_{i, j}$ is a partition derived from the Cauchy-Binet expansion for each Δ 's, s_μ is a Schur function.

Prop 1. The map $f : x \mapsto \tilde{P}_R(x)$ is a homomorphism on a “proper” subset $S \subseteq \mathcal{P}$, s.t. for any $x, y \in S$

$$\tilde{P}_R(x \cdot y) = \frac{1}{2} \sum_{\mu, \eta, \nu} \left(\text{coeff}(\tilde{P}_R(x), \Delta_\mu) - \text{coeff}(\tilde{P}_R(y), \Delta_\eta) \right) \Delta_\nu \simeq_{|Res(\Delta)} \frac{1}{2} \left(\tilde{P}_R(x) - \tilde{P}_R(y) \right) \quad (49)$$

Note: this construction is very sensitive to the choice of R (if we swap the transposition matrices, S will change).

Lemma 2. If $x \in \mathcal{P}(Q)$ is of spiral type 1, then


$$\tilde{P}_\pm(x) \equiv 0, \quad (50)$$

[illegible]

Another (stand-alone) example:

$$-IP_6(\text{spiral}_2) = s_{\begin{array}{|c|} \hline \square \\ \square \\ \hline \end{array}_2} + s_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}_0} + s_{\begin{array}{|c|} \hline \square \\ \square & \square \\ \hline \end{array}_4} - s_{\begin{array}{|c|} \hline \square \\ \square & \square \\ \hline \end{array}_3} - s_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}_5} - s_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}_1} \quad (53)$$

Let $(m, n) = (6, 6)$; consider the following two paths and their concatenation:



$$\begin{aligned}
\tilde{P}(Q_1) &= \left(s_{\square_0} + s_{\square_3} + s_{\square_5} + s_{\square_2} \right) \Delta_{\square_5} + \left(-s_{\square_0} + s_{\square_3} + s_{\square_5} - s_{\square_2} \right) \Delta_{\square_5} + \\
&\quad + \left(s_{\square_2} + s_{\square_0} + s_{\square_5} + s_{\square_3} \right) \Delta_{\square_5} + \left(s_{\square_5} + s_{\square_0} + s_{\square_3} + s_{\square_2} \right) \Delta_{\square_5} + \\
&\quad + \left(-s_{\square_2} + s_{\square_3} + s_{\square_5} - s_{\square_0} \right) \Delta_{\square_5} + \left(s_{\square_0} + s_{\square_2} + s_{\square_5} + s_{\square_3} \right) \Delta_{\square_5} + \\
&\quad + \left(s_{\square_5} + s_{\square_2} + s_{\square_0} + s_{\square_3} \right) \Delta_{\square_5} + \left(s_{\square_3} - s_{\square_2} + s_{\square_5} - s_{\square_0} \right) \Delta_{\square_5} + \\
&\quad + \left(s_{\square_5} + s_{\square_3} + s_{\square_2} + s_{\square_0} \right) \Delta_{\square_5} \quad \# 5 \text{ is the longest arc} \\
\tilde{P}(Q_2) &= \left(s_{\square_4} + s_{\square_5} - s_{\square_2} + s_{\square_1} \right) \Delta_{\square_4} + \left(-s_{\square_5} - s_{\square_1} + s_{\square_2} + s_{\square_4} \right) \Delta_{\square_4} + \\
&\quad + \left(-s_{\square_5} + s_{\square_2} + s_{\square_4} + s_{\square_1} \right) \Delta_{\square_4} + \left(-s_{\square_2} + s_{\square_1} + s_{\square_4} + s_{\square_5} \right) \Delta_{\square_4} + \\
&\quad + \left(-s_{\square_5} + s_{\square_2} + s_{\square_4} - s_{\square_1} \right) \Delta_{\square_4} + \left(s_{\square_2} - s_{\square_5} + s_{\square_1} + s_{\square_4} \right) \Delta_{\square_4} + \\
&\quad + \left(s_{\square_1} + s_{\square_5} + s_{\square_4} - s_{\square_2} \right) \Delta_{\square_4} + \left(-s_{\square_1} - s_{\square_5} + s_{\square_4} + s_{\square_2} \right) \Delta_{\square_4} + \\
&\quad + \left(s_{\square_2} + s_{\square_1} + s_{\square_4} - s_{\square_5} \right) \Delta_{\square_4} \quad \# 4 \text{ is the longest arc} \\
\tilde{P}(Q_{1,2}) &= \left(-s_{\square_2} + s_{\square_1} - s_{\square_4} - s_{\square_0} + s_{\square_3} + s_{\square_5} \right) \Delta_{\square_3} + \left(s_{\square_5} + s_{\square_1} - s_{\square_2} - s_{\square_4} \right) \Delta_{\square_3} + \\
&\quad + \left(s_{\square_5} - s_{\square_0} - s_{\square_4} + s_{\square_3} + s_{\square_1} - s_{\square_2} \right) \Delta_{\square_3} = \\
&= \frac{1}{2} \left(\text{coeff}(P(Q_1), \Delta_{\square_5}) - \text{coeff}(P(Q_2), \Delta_{\square_4}) \right) \Delta_{\square_3} + \frac{1}{2} \left(\text{coeff}(P(Q_1), \Delta_{\square_5}) - \text{coeff}(P(Q_2), \Delta_{\square_4}) \right) \Delta_{\square_3} + \\
&\quad + \frac{1}{2} \left(\text{coeff}(P(Q_1), \Delta_{\square_5}) - \text{coeff}(P(Q_2), \Delta_{\square_4}) \right) \Delta_{\square_3} = \frac{1}{2} (P(Q_1) - \text{permut}(P(Q_2), \Delta_*))
\end{aligned}$$

(This is a particular example related to the proposition 1).

The next goal is to build infinite series like this

$$Z_R(S) = \frac{1}{2k} \sum_{i,j} \left(\tilde{P}_R(x_i) - \tilde{P}_R(x_j) \right), \quad (55)$$

where $S = \{x_1, x_2, \dots, x_i, \dots\} \subseteq \mathcal{P}(Q_{n \rightarrow \infty})$ is a proper union of loops. Of course it can be rewritten in




Figure 4: Concatenation of $x_i \in S$ for $R = \Sigma_0$

terms of Schur function expansion

$$Z_R(S) = \sum_{\lambda} g_{\lambda}(\Delta) s_{\lambda}(p_1, p_2, \dots), \quad (56)$$

or, in our “ladder notation”, over all Γ -type *weighted* Young diagrams, using the $\mathcal{Y}$ operator

$$Z_R(S) = \sum_{\lambda_w} g_{\lambda_w}(\Delta) \tilde{s}_{\lambda_w}(p_1, p_2, \dots), \quad w \in \mathbb{N}, \quad \lambda_w = \mathcal{Y}(\lambda) \quad (57)$$

Take $S' = \{x_1, x_2\} \subset \mathcal{P}(Q_{10})$, $(m, n) = (10, 10)$. In this case the double-path polynomial is non-trivial (!)

$$\begin{aligned} \tilde{P}_{\pm}(S') = & \left(s_{\square_2} + s_{\begin{array}{|c|c|c|c|c|} \hline \square & \square & \square & \square & \square \\ \hline \end{array}} + s_{\square_6} + s_{\square_3} + s_{\begin{array}{|c|c|c|c|c|} \hline \square & \square & \square & \square & \square \\ \hline \end{array}} + s_{\square_1} + s_{\square_7} + s_{\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}} + s_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} \right) \Delta_{\square_6} + \\ & + \left(s_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} + s_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} + s_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} + s_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} + s_{\begin{array}{|c|c|c|c|c|} \hline \square & \square & \square & \square & \square \\ \hline \end{array}} + s_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} + s_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} + s_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} + s_{\begin{array}{|c|c|c|c|c|} \hline \square & \square & \square & \square & \square \\ \hline \end{array}} \right) \Delta_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} + \\ & + \left(s_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} + s_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} + s_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} + s_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} + s_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} + s_{\begin{array}{|c|c|c|c|c|} \hline \square & \square & \square & \square & \square \\ \hline \end{array}} + s_{\begin{array}{|c|c|c|c|c|} \hline \square & \square & \square & \square & \square \\ \hline \end{array}} + s_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} + s_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} \right) \Delta_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} + \\ & + \left(s_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} + s_{\begin{array}{|c|c|c|c|c|} \hline \square & \square & \square & \square & \square \\ \hline \end{array}} + s_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} + s_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} + s_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} + s_{\begin{array}{|c|c|c|c|c|} \hline \square & \square & \square & \square & \square \\ \hline \end{array}} + s_{\begin{array}{|c|c|c|c|c|} \hline \square & \square & \square & \square & \square \\ \hline \end{array}} + s_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} + s_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} \right) \Delta_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} + \\ & + s_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} + s_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} + s_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} + s_{\begin{array}{|c|c|c|c|c|} \hline \square & \square & \square & \square & \square \\ \hline \end{array}} + s_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} + s_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} + s_{\begin{array}{|c|c|c|c|c|} \hline \square & \square & \square & \square & \square \\ \hline \end{array}} + s_{\begin{array}{|c|} \hline \square \\ \hline \end{array}} \end{aligned} \quad (58)$$

(Here we assumed $\Delta_{[6,7,8,9,10,16,17,18,19,20,\dots,96,97,98,99,100]} = 1$ to simplify the formula). Its configuration is drawn on the figure 5 (right). Now let's extend the picture for x_3 : The problem is that in this case $(m, n) = (16, 16)$ the image is empty, so we should add some isolated points to the graph 5 (left).





Figure 5: Concatenation of $x_1, x_2 \in S'$ for $R = \Sigma_0$: compatible case

Let

$$X = \{G_{i_1}^{(j_1)}, G_{i_2}^{(j_2)}, \dots, G_{i_s}^{(j_s)}\}^k \in \mathcal{P} \quad (59)$$

be an element of path algebra of $Q_{\text{univ}}(k)$, built as concatenation normal loops G_{i_1}, G_{i_2} , each one shifted on the j -th position. For example, consider 3-union $\{G_4^{(0)}, G_6^{(3)}, G_8^{(2)}\}^{10}$. QR has unique solution; furthermore, we take $\Delta_{\begin{smallmatrix} \square \\ \square \\ 7 \end{smallmatrix}} = 1$ (denominator) to homogenise the resulting formula. Here is the associated quiver:



Then $\tilde{P}_{\pm}(X)$ become homogeneous in Δ :

$$\begin{aligned} \tilde{P}_{\pm}(X) = & \left(s_{\begin{smallmatrix} \square \\ \square \\ \square \\ \square \\ 9 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 0 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 5 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 1 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 2 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 4 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 6 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 3 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ \square \\ \square \\ 8 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 7 \end{smallmatrix}} \right) \Delta_{\begin{smallmatrix} \square \\ \square \\ 6 \end{smallmatrix}} + \\ & + \left(s_{\begin{smallmatrix} \square \\ \square \\ 0 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ \square \\ \square \\ 8 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 3 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 1 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 6 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 5 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 2 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 4 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 7 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ \square \\ \square \\ 9 \end{smallmatrix}} \right) \Delta_{\begin{smallmatrix} \square \\ \square \\ 6 \end{smallmatrix}} + \\ & + \left(s_{\begin{smallmatrix} \square \\ \square \\ 2 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 7 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 6 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 3 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 5 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 4 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ \square \\ \square \\ 9 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 0 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 1 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ \square \\ \square \\ 8 \end{smallmatrix}} \right) \Delta_{\begin{smallmatrix} \square \\ \square \\ 6 \end{smallmatrix}} + \\ & + \left(s_{\begin{smallmatrix} \square \\ \square \\ \square \\ \square \\ 9 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ \square \\ \square \\ 8 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 1 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 7 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 5 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 6 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 0 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 2 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 3 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 4 \end{smallmatrix}} \right) \Delta_{\begin{smallmatrix} \square \\ \square \\ 6 \end{smallmatrix}} + \\ & + s_{\begin{smallmatrix} \square \\ \square \\ 0 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 1 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ \square \\ \square \\ 8 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 5 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 3 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ \square \\ \square \\ 9 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 4 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 6 \end{smallmatrix}} + s_{\begin{smallmatrix} \square \\ \square \\ 2 \end{smallmatrix}} \end{aligned} \quad (61)$$

Remark: we can see only antisymmetric coefficients Δ here! To extend this case for G_{10} one should exclude $G_8^{(2)}$.




Figure 6: Non-trivial concatenation of loops $X = \{G_4^{(0)}, G_6^{(3)}, G_8^{(2)}\}^{10}$ and $X' = \{G_4^{(0)}, G_6^{(3)}, G_8^{(2)}, \underline{G_{10}^{(2)}}\}^{10}$ (only the largest arcs are shown)

The resulting picture is that each loop can be scaled separately, such that increasing k, s simultaneously does not change the consistency of QR. Our next aim is to investigate asymptotical properties, derived from $\{G_{i_1}^{(j_1)}, G_{i_2}^{(j_2)}, \dots, G_{i_s}^{(j_s)}\}^k$ when $k, s \rightarrow \infty$. This is of further development.

References

- [1] ALISTAIR SAVAGE, PETER TINGLEY Quiver grassmannians, quiver varieties and the preprojective algebra // <http://arxiv.org/abs/0909.3746>
- [2] MAXIM KONTSEVICH, YAN SOIBELMAN Lectures on motivic Donaldson-Thomas invariants and wall-crossing formulas // arxiv preprint, 2011
- [3] OLIVER LORSCHIED, THORSTEN WEIST Quiver Grassmannians of type extended Dynkin type D - Part 1: Schubert systems and decompositions into affine spaces // <https://arxiv.org/abs/1507.00392>
- [4] OLIVER LORSCHIED, THORSTEN WEIST Homogeneous coordinates for quiver grassmannians // <http://arxiv.org/abs/1607.01058v1>
- [5] OLIVER LORSCHIED, THORSTEN WEIST Quiver Grassmannians of extended Dynkin type D - Part 2: Schubert decompositions and F-polynomials // <http://arxiv.org/abs/1507.00395>